



**MONJUVI**<sup>®</sup>  
tafasitamab-cxix | 200mg  
for injection, for intravenous use

# PRESCRIBING INFORMATION

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MONJUVI safely and effectively. See full prescribing information for MONJUVI.

MONJUVI<sup>®</sup> (tafasitamab-cxix) for injection, for intravenous use  
Initial U.S. Approval: 2020

## INDICATIONS AND USAGE

MONJUVI is a CD19-directed cytolytic antibody indicated in combination with lenalidomide for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma and who are not eligible for autologous stem cell transplant (ASCT).

This indication is approved under accelerated approval based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s). (1)

## DOSAGE AND ADMINISTRATION

- Administer premedications prior to starting MONJUVI. (2.2)
- The recommended dosage of MONJUVI is 12 mg/kg as an intravenous infusion according to the following dosing schedule: (2.1)
  - Cycle 1: Days 1, 4, 8, 15 and 22 of the 28-day cycle.
  - Cycles 2 and 3: Days 1, 8, 15 and 22 of each 28-day cycle.
  - Cycle 4 and beyond: Days 1 and 15 of each 28-day cycle.
- Administer MONJUVI in combination with lenalidomide for a maximum of 12 cycles and then continue MONJUVI as monotherapy until disease progression or unacceptable toxicity. (2.1)
- See Full Prescribing Information for instructions on preparation and administration. (2.3, 2.4)

## DOSAGE FORMS AND STRENGTHS

For injection: 200 mg of tafasitamab-cxix as lyophilized powder in single-dose vial for reconstitution. (3)

## CONTRAINDICATIONS

None.

## WARNINGS AND PRECAUTIONS

- Infusion-Related Reactions: Monitor patients frequently during infusion. Interrupt or discontinue infusion based on severity. (2.3, 5.1)
- Myelosuppression: Monitor complete blood counts. Manage using dose modifications and growth factor support. Interrupt or discontinue MONJUVI based on severity. (2.3, 5.2)
- Infections: Bacterial, fungal and viral infections can occur during and following MONJUVI. Monitor patients for infections. (2.3, 5.3)
- Embryo-Fetal Toxicity: May cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and use of effective contraception. (5.4)

## ADVERSE REACTIONS

The most common adverse reactions ( $\geq 20\%$ ) are neutropenia, fatigue, anemia, diarrhea, thrombocytopenia, cough, pyrexia, peripheral edema, respiratory tract infection, and decreased appetite. (6.1)

To report **SUSPECTED ADVERSE REACTIONS**, contact MORPHOSYS US INC. at 1-844-667-1992 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## USE IN SPECIFIC POPULATIONS

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling  
Revised: 7/2020

# FULL PRESCRIBING INFORMATION: CONTENTS\*

## 1 INDICATIONS AND USAGE

## 2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Recommended Premedications
- 2.3 Dosage Modifications for Adverse Reactions
- 2.4 Preparation and Administration

## 3 DOSAGE FORMS AND STRENGTHS

## 4 CONTRAINDICATIONS

## 5 WARNINGS AND PRECAUTIONS

- 5.1 Infusion-Related Reactions
- 5.2 Myelosuppression
- 5.3 Infections
- 5.4 Embryo-Fetal Toxicity

## 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Immunogenicity

## 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

## 11 DESCRIPTION

## 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

## 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## 14 CLINICAL STUDIES

## 16 HOW SUPPLIED/STORAGE AND HANDLING

## 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

# FULL PRESCRIBING INFORMATION

## 1 INDICATIONS AND USAGE

MONJUVI, in combination with lenalidomide, is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant (ASCT).

This indication is approved under accelerated approval based on overall response rate [see *Clinical Studies (14)*]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Recommended Dosage

The recommended dose of MONJUVI is 12 mg/kg based on actual body weight administered as an intravenous infusion according to the dosing schedule in Table 1.

Administer MONJUVI in combination with lenalidomide 25 mg for a maximum of 12 cycles, then continue MONJUVI as monotherapy until disease progression or unacceptable toxicity [see *Clinical Studies (14)*]. Refer to the lenalidomide prescribing information for lenalidomide dosage recommendations.

**TABLE 1: MONJUVI Dosing Schedule**

Cycle*	Dosing schedule
<b>Cycle 1</b>	Days 1, 4, 8, 15 and 22
<b>Cycles 2 and 3</b>	Days 1, 8, 15 and 22
<b>Cycle 4 and beyond</b>	Days 1 and 15

\*Each treatment cycle is 28-days.

MONJUVI should be administered by a healthcare professional with immediate access to emergency equipment and appropriate medical support to manage infusion-related reactions (IRRs) [see *Warnings and Precautions (5.1)*].

### 2.2 Recommended Premedications

Administer premedications 30 minutes to 2 hours prior to starting MONJUVI infusion to minimize infusion-related reactions [see *Warning and Precautions (5.1)*]. Premedications may include acetaminophen, histamine H<sub>1</sub> receptor antagonists, histamine H<sub>2</sub> receptor antagonists, and/or glucocorticosteroids.

## 2.2 Recommended Premedications (continued)

For patients not experiencing infusion-related reactions during the first 3 infusions, premedication is optional for subsequent infusions.

If a patient experiences an infusion-related reaction, administer premedications before each subsequent infusion.

## 2.3 Dosage Modifications for Adverse Reactions

The recommended dosage modifications for adverse reactions are summarized in Table 2.

**TABLE 2: Dosage Modifications for Adverse Reactions**

Adverse Reaction	Severity	Dosage Modification
Infusion-related reactions <i>[see Warnings and Precautions (5.1)]</i>	Grade 2 (moderate)	<ul style="list-style-type: none"> <li>Interrupt infusion immediately and manage signs and symptoms.</li> <li>Once signs and symptoms resolve or reduce to Grade 1, resume infusion at no more than 50% of the rate at which the reaction occurred. If the patient does not experience further reaction within 1 hour and vital signs are stable, the infusion rate may be increased every 30 minutes as tolerated to the rate at which the reaction occurred.</li> </ul>
	Grade 3 (severe)	<ul style="list-style-type: none"> <li>Interrupt infusion immediately and manage signs and symptoms.</li> <li>Once signs and symptoms resolve or reduce to Grade 1, resume infusion at no more than 25% of the rate at which the reaction occurred. If the patient does not experience further reaction within 1 hour and vital signs are stable, the infusion rate may be increased every 30 minutes as tolerated to a maximum of 50% of the rate at which the reaction occurred.</li> <li>If after rechallenge the reaction returns, stop the infusion immediately.</li> </ul>
	Grade 4 (life-threatening)	<ul style="list-style-type: none"> <li>Stop the infusion immediately and permanently discontinue MONJUVI.</li> </ul>

Adverse Reaction	Severity	Dosage Modification
<b>Myelosuppression</b> <i>[see Warnings and Precautions (5.2)]</i>	Platelet count of 50,000/ mL or less	<ul style="list-style-type: none"> <li>Withhold MONJUVI and lenalidomide and monitor complete blood count (CBC) weekly until platelet count is 50,000/ mL or higher.</li> <li>Resume MONJUVI at the same dose and lenalidomide at a reduced dose. Refer to lenalidomide prescribing information for dosage modifications.</li> </ul>
	Neutrophil count of 1,000/ mL or less for at least 7 days OR Neutrophil count of 1,000/ mL or less with an increase of body temperature to 100.4°F (38°C) or higher OR Neutrophil count less than 500/mL	<ul style="list-style-type: none"> <li>Withhold MONJUVI and lenalidomide and monitor CBC weekly until neutrophil count is 1,000/ mL or higher.</li> <li>Resume MONJUVI at the same dose and lenalidomide at a reduced dose. Refer to lenalidomide prescribing information for dosage modifications.</li> </ul>

## 2.4 Preparation and Administration

Reconstitute and dilute MONJUVI prior to infusion.

### Reconstitution

1. Calculate the dose (mg) and determine the number of vials needed.
2. Reconstitute each 200 mg MONJUVI vial with 5 mL Sterile Water for Injection, USP with the stream directed toward the wall of each vial to obtain a final concentration of 40 mg/mL tafasitamab-cxix.
3. Gently swirl the vial(s) until completely dissolved. Do not shake or swirl vigorously. Complete dissolution may take up to 5 minutes.
4. Visually inspect the reconstituted solution for particulate matter or discoloration. The reconstituted solution should appear as a colorless to slightly yellow solution. Discard the vial(s) if the solution is cloudy, discolored, or contains visible particles.
5. Use the reconstituted MONJUVI solution immediately. If needed, store the reconstituted solution in the vial for a maximum of 12 hours either refrigerated at 36°F to 46°F (2°C to 8°C) or room temperature at 68°F to 77°F (20°C to 25°C) before dilution. Protect from light during storage.

## 2.4 Preparation and Administration (continued)

### Dilution

1. Determine the volume (mL) of the 40 mg/mL reconstituted MONJUVI solution needed based on the required dose.
2. Remove a volume equal to the required MONJUVI solution from a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag and discard it.
3. Withdraw the necessary amount of MONJUVI and slowly dilute in the infusion bag that contains the 0.9% Sodium Chloride Injection, USP to a final concentration of 2 mg/mL to 8 mg/mL. Discard any unused portion of MONJUVI remaining in the vial.
4. Gently mix the intravenous bag by slowly inverting the bag. Do not shake. Visually inspect the infusion bag with the diluted MONJUVI infusion solution for particulate matter and discoloration prior to administration.
5. If not used immediately, store the diluted MONJUVI infusion solution refrigerated for up to 18 hours at 36°F to 46°F (2°C to 8°C) and/or at room temperature for up to 12 hours at 68°F to 77°F (20°C to 25°C). The room temperature storage includes time for infusion. Protect from light during storage.

Do not shake or freeze the reconstituted or diluted infusion solutions.

### Administration

- Administer MONJUVI as an intravenous infusion.
  - For the first infusion, use an infusion rate of 70 mL/h for the first 30 minutes, then, increase the rate so that the infusion is administered within 1.5 to 2.5 hours.
  - Administer all subsequent infusions within 1.5 to 2 hours.
- Infuse the entire contents of the bag containing MONJUVI.
- Do not co-administer other drugs through the same infusion line.
- No incompatibilities have been observed between MONJUVI with infusion containers made of polypropylene (PP), polyvinylchloride (PVC), polyethylene (PE), polyethyleneterephthalate (PET), or glass and infusion sets made of polyurethane (PUR) or PVC.

## 3 DOSAGE FORMS AND STRENGTHS

For injection: 200 mg of tafasitamab-cxix as white to slightly yellowish lyophilized powder in single dose vial for reconstitution and further dilution.

## 4 CONTRAINDICATIONS

None.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Infusion-Related Reactions

MONJUVI can cause infusion-related reactions [see *Adverse Reactions (6.1)*]. In L-MIND, infusion-related reactions occurred in 6% of the 81 patients. Eighty percent of infusion-related reactions occurred during cycle 1 or 2. Signs and symptoms included chills, flushing, dyspnea, and hypertension. These reactions were managed with temporary interruption of the infusion and/or with supportive medication.

Premedicate patients prior to starting MONJUVI infusion [see *Dosage and Administration (2.2)*]. Monitor patients frequently during infusion. Based on the severity of the infusion-related reaction, interrupt or discontinue MONJUVI [see *Dosage and Administration (2.3)*]. Institute appropriate medical management.

### 5.2 Myelosuppression

MONJUVI can cause serious or severe myelosuppression, including neutropenia, thrombocytopenia, and anemia [see *Adverse Reactions (6.1)*]. In L-MIND, Grade 3 neutropenia occurred in 25% of patients, thrombocytopenia in 12% and anemia in 7%. Grade 4 neutropenia occurred in 25% and thrombocytopenia in 6%. Neutropenia led to treatment discontinuation in 3.7% of patients.

Monitor CBC prior to administration of each treatment cycle and throughout treatment. Monitor patients with neutropenia for signs of infection. Consider granulocyte colony-stimulating factor administration. Withhold MONJUVI based on the severity of the adverse reaction [see *Dosage and Administration (2.3)*]. Refer to the lenalidomide prescribing information for dosage modifications.

### 5.3 Infections

Fatal and serious infections, including opportunistic infections, occurred in patients during treatment with MONJUVI and following the last dose [see *Adverse Reactions (6.1)*].

In L-MIND, 73% of the 81 patients developed an infection. The most frequent infections were respiratory tract infection (24%), urinary tract infection (17%), bronchitis (16%), nasopharyngitis (10%) and pneumonia (10%). Grade 3 or higher infection occurred in 30% of the 81 patients. The most frequent grade 3 or higher infection was pneumonia (7%). Infection-related deaths were reported in 2.5% of the 81 patients.

Monitor patients for signs and symptoms of infection and manage infections as appropriate.

### 5.4 Embryo-Fetal Toxicity

Based on its mechanism of action, MONJUVI may cause fetal B-cell depletion when administered to a pregnant woman. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with MONJUVI and for at least 3 months after the last dose [see *Use in Specific Populations (8.1, 8.3)*].

MONJUVI is initially administered in combination with lenalidomide. The combination of MONJUVI with lenalidomide is contraindicated in pregnant women because lenalidomide can cause birth defects and death of the unborn child. Refer to the lenalidomide prescribing information on use during pregnancy.

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Infusion-related reactions [see *Warnings and Precautions (5.1)*]
- Myelosuppression [see *Warnings and Precautions (5.2)*]
- Infections [see *Warnings and Precautions (5.3)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in other clinical trials of another drug and may not reflect the rates observed in practice.

#### Relapsed or Refractory Diffuse Large B-Cell Lymphoma

The safety of MONJUVI was evaluated in L-MIND [see *Clinical Studies (14)*]. Patients (n=81) received MONJUVI 12 mg/kg intravenously in combination with lenalidomide for maximum of 12 cycles, followed by MONJUVI as monotherapy until disease progression or unacceptable toxicity as follows:

- Cycle 1: Days 1, 4, 8, 15 and 22 of the 28-day cycle;
- Cycle 2 and 3: Days 1, 8, 15 and 22 of each 28-day cycle;
- Cycles 4 and beyond: Days 1 and 15 of each 28-day cycle.

Among patients who received MONJUVI, 57% were exposed for 6 months or longer, 42% were exposed for greater than one year, and 24% were exposed for greater than two years.

Serious adverse reactions occurred in 52% of patients who received MONJUVI. Serious adverse reactions in  $\geq 6\%$  of patients included infections (26%), including pneumonia (7%), and febrile neutropenia (6%). Fatal adverse reactions occurred in 5% of patients who received MONJUVI, including cerebrovascular accident (1.2%), respiratory failure (1.2%), progressive multifocal leukoencephalopathy (1.2%) and sudden death (1.2%).

Permanent discontinuation of MONJUVI or lenalidomide due to an adverse reaction occurred in 25% of patients and permanent discontinuation of MONJUVI due to an adverse reaction occurred in 15%. The most frequent adverse reactions which resulted in permanent discontinuation of MONJUVI were infections (5%), nervous system disorders (2.5%), respiratory, thoracic and mediastinal disorders (2.5%).

Dosage interruptions of MONJUVI or lenalidomide due to an adverse reaction occurred in 69% of patients and dosage interruption of MONJUVI due to an adverse reaction occurred in 65%. The most frequent adverse reactions which required a dosage interruption of MONJUVI were blood and lymphatic system disorders (41%), and infections (27%).

The most common adverse reactions ( $\geq 20\%$ ) were neutropenia, fatigue, anemia, diarrhea, thrombocytopenia, cough, pyrexia, peripheral edema, respiratory tract infection, and decreased appetite.

Table 3 summarizes the adverse reactions in L-MIND.

**TABLE 3: Adverse Reactions (≥10%) in Patients with Relapsed or Refractory Diffuse Large B-Cell Lymphoma Who Received MONJUVI in L-MIND**

Adverse Reaction	MONJUVI (N=81)	
	All Grades (%)	Grade 3 or 4 (%)
<b>Blood and lymphatic system disorders</b>		
Neutropenia	51	49
Anemia	36	7
Thrombocytopenia	31	17
Febrile neutropenia	12	12
<b>General disorders and administration site conditions</b>		
Fatigue*	38	3.7
Pyrexia	24	1.2
Peripheral edema	24	0
<b>Gastrointestinal disorders</b>		
Diarrhea	36	1.2
Constipation	17	0
Nausea	15	0
Vomiting	15	0
<b>Respiratory, thoracic and mediastinal disorders</b>		
Cough	26	1.2
Dyspnea	12	1.2

## 6.1 Clinical Trials Experience (continued)

Adverse Reaction	MONJUVI (N=81)	
	All Grades (%)	Grade 3 or 4 (%)
<b>Infections</b>		
Respiratory tract infection <sup>†</sup>	24	4.9
Urinary tract infection <sup>†</sup>	17	4.9
Bronchitis	16	1.2
<b>Metabolism and nutrition disorders</b>		
Decreased appetite	22	0
Hypokalemia	19	6
<b>Musculoskeletal and connective tissue disorders</b>		
Back pain	19	2.5
Muscle spasms	15	0

\* Fatigue includes asthenia and fatigue

<sup>†</sup> Respiratory tract infection includes: lower respiratory tract infection, upper respiratory tract infection, respiratory tract infection

<sup>†</sup> Urinary tract infection includes: urinary tract infection, Escherichia urinary tract infection, urinary tract infection bacterial, urinary tract infection enterococcal

Clinically relevant adverse reactions in <10% of patients who received MONJUVI were:

- *Blood and lymphatic system disorders*: lymphopenia (6%)
- *General disorders and administration site conditions*: infusion-related reaction (6%)
- *Infections*: sepsis (4.9%)
- *Investigations*: weight decreased (4.9%)
- *Musculoskeletal and connective tissue disorders*: arthralgia (9%), pain in extremity (9%), musculoskeletal pain (2.5%)
- *Neoplasms benign, malignant and unspecified*: basal cell carcinoma (1.2%)
- *Nervous system disorders*: headache (9%), paresthesia (7%), dysgeusia (6%)
- *Respiratory, thoracic and mediastinal disorders*: nasal congestion (4.9%), exacerbation of chronic obstructive pulmonary disease (1.2%)
- *Skin and subcutaneous tissue disorders*: erythema (4.9%), alopecia (2.5%), hyperhidrosis (2.5%)

Table 4 summarizes the laboratory abnormalities in L-MIND.

**TABLE 4: Select Laboratory Abnormalities (>20%) Worsening from Baseline in Patients with Relapsed or Refractory Diffuse Large B-Cell Lymphoma Who Received MONJUVI in L-MIND**

Laboratory Abnormality	MONJUVI <sup>1</sup>	
	All Grades (%)	Grade 3 or 4 (%)
<b>Chemistry</b>		
Glucose increased	49	5
Calcium decreased	47	1.4
Gamma glutamyl transferase increased	34	5
Albumin decreased	26	0
Magnesium decreased	22	0
Urate increased	20	7
Phosphate decreased	20	5
Creatinine increased	20	1.4
Aspartate aminotransferase increased	20	0
<b>Coagulation</b>		
Activated partial thromboplastin time increased	46	4.1

<sup>1</sup>The denominator used to calculate the rate was 74 based on the number of patients with a baseline value and at least one post-treatment value.

## 6.2 Immunogenicity

As with all therapeutic proteins, there is the potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assays. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors, including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies in the studies described below with the incidence of antibodies in other studies or to other tafasitamab products may be misleading.

Overall, no treatment-emergent or treatment-boosted anti-tafasitamab antibodies were observed. No clinically meaningful differences in the pharmacokinetics, efficacy, or safety profile of tafasitamab-cxix were observed in 2.5% of 81 patients with relapsed or refractory DLBCL with pre-existing anti-tafasitamab antibodies in L-MIND.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Based on its mechanism of action, MONJUVI may cause fetal B-cell depletion when administered to a pregnant woman [see *Clinical Pharmacology (12.1)*]. There are no available data on MONJUVI use in pregnant women to evaluate for a drug-associated risk. Animal reproductive toxicity studies have not been conducted with tafasitamab-cxix.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

MONJUVI is administered in combination with lenalidomide for up to 12 cycles. Lenalidomide can cause embryo-fetal harm and is contraindicated for use in pregnancy. Refer to the lenalidomide prescribing information for additional information. Lenalidomide is only available through a REMS program.

#### Clinical Considerations

##### *Fetal/Neonatal Adverse Reactions*

Immunoglobulin G (IgG) monoclonal antibodies are transferred across the placenta. Based on its mechanism of action, MONJUVI may cause depletion of fetal CD19 positive immune cells. Defer administering live vaccines to neonates and infants exposed to tafasitamab-cxix in utero until a hematology evaluation is completed.

#### Data

##### *Animal Data*

Animal reproductive studies have not been conducted with tafasitamab-cxix. Tafasitamab-cxix is an IgG antibody and thus has the potential to cross the placental barrier permitting direct fetal exposure and depleting fetal B lymphocytes.

### 8.2 Lactation

#### Risk Summary

There are no data on the presence of tafasitamab-cxix in human milk or the effects on the breastfed child or milk production. Maternal immunoglobulin G is known to be present in human milk. The effects of local gastrointestinal exposure and limited systemic exposure in the breastfed infant to MONJUVI are unknown. Because of the potential for serious adverse reactions in the breastfed child, advise women not to breastfeed during treatment with MONJUVI and for at least 3 months after the last dose. Refer to lenalidomide prescribing information for additional information.

## 8.3 Females and Males of Reproductive Potential

MONJUVI can cause fetal B-cell depletion when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

### Pregnancy Testing

Refer to the prescribing information for lenalidomide for pregnancy testing requirements prior to initiating the combination of MONJUVI with lenalidomide.

### Contraception

#### *Females*

Advise females of reproductive potential to use effective contraception during treatment with MONJUVI and for at least 3 months after the last dose. Additionally, refer to the lenalidomide prescribing information for additional recommendations for contraception.

#### *Males*

Refer to the lenalidomide prescribing information for recommendations.

## 8.4 Pediatric Use

The safety and effectiveness of MONJUVI in pediatric patients have not been established.

## 8.5 Geriatric Use

Among 81 patients who received MONJUVI and lenalidomide in L-MIND, 72% were 65 years and older, while 38% were 75 years and older. Clinical studies of MONJUVI did not include sufficient numbers of patients aged 65 and older to determine whether effectiveness differs compared to that of younger subjects. Patients 65 years and older had more serious adverse reactions (57%) than younger patients (39%).

## 11 DESCRIPTION

Tafasitamab-cxix is a humanized CD19-directed cytolytic monoclonal antibody that contains an IgG1/2 hybrid Fc-domain with 2 amino acid substitutions to modify the Fc-mediated functions of the antibody. It is produced by recombinant DNA technology in mammalian cells (Chinese hamster ovary). Tafasitamab-cxix has a molecular weight of approximately 150 kDa.

MONJUVI (tafasitamab-cxix) for injection is supplied as a sterile, preservative-free, white to slightly yellowish lyophilized powder in a single-dose vial for intravenous use after reconstitution and further dilution. After reconstitution with 5 mL of Sterile Water for Injection, USP, the resulting concentration is 40 mg/mL with a pH of 6.0. Each single-dose vial contains 200 mg tafasitamab-cxix, citric acid monohydrate (3.7 mg), polysorbate 20 (1 mg), sodium citrate dihydrate (31.6 mg) and trehalose dihydrate (378.3 mg).

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Tafasitamab-cxix is an Fc-modified monoclonal antibody that binds to CD19 antigen expressed on the surface of pre-B and mature B lymphocytes and on several B-cell malignancies, including diffuse large B-cell lymphoma (DLBCL).

Upon binding to CD19, tafasitamab-cxix mediates B-cell lysis through apoptosis and immune effector mechanisms, including antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP).

In studies conducted in vitro in DLBCL tumor cells, tafasitamab-cxix in combination with lenalidomide resulted in increased ADCC activity compared to tafasitamab-cxix or lenalidomide alone.

### 12.2 Pharmacodynamics

Tafasitamab-cxix reduced peripheral blood B cell counts by 97% after eight days of treatment in patients with relapsed or refractory DLBCL. Nadir, with a reduction of 100%, was reached within 16 weeks of treatment.

### 12.3 Pharmacokinetics

Mean trough concentrations ( $\pm$  standard deviation) were 179 ( $\pm$  53)  $\mu\text{g/mL}$  following administration of MONJUVI at 12 mg/kg on Days 1, 8, 15, and 22 in Cycle 1-3 (plus an additional dose on Cycle 1 Day 4), and 153 ( $\pm$  68)  $\mu\text{g/mL}$  following administration of MONJUVI at 12 mg/kg on Days 1 and 15 from Cycle 4 onwards. Overall maximum tafasitamab-cxix serum concentrations were 483 ( $\pm$ 109)  $\mu\text{g/mL}$ .

#### Distribution

The total volume of distribution for tafasitamab-cxix was 9.3 L (95% CI: 8.6, 10 L).

#### Elimination

The clearance of tafasitamab-cxix was 0.41 L/day (CV: 32%) and terminal elimination half-life was 17 days (95% CI: 15, 18 days).

#### Specific Populations

Bodyweight (40 to 163 kg) has a significant effect on the pharmacokinetics of tafasitamab-cxix, with higher clearance and volume of distribution expected with higher body weight. No clinically meaningful differences in the pharmacokinetics of tafasitamab-cxix were observed based on age (16 to 90 years), sex, mild to moderate renal impairment (CLcr 30-89 mL/min estimated by the Cockcroft-Gault equation), and mild hepatic impairment (total bilirubin  $\leq$  ULN and AST  $>$  ULN, or total bilirubin 1 to 1.5 times ULN and any AST). The effect of severe renal impairment to end-stage renal disease (CLcr  $<$  30 mL/min), moderate to severe hepatic impairment (total bilirubin  $>$  1.5 times ULN and any AST), and race/ethnicity on tafasitamab-cxix pharmacokinetics is unknown.

#### Drug Interaction Studies

No clinically meaningful differences in tafasitamab-cxix pharmacokinetics were observed when used concomitantly with lenalidomide.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity and genotoxicity studies have not been conducted with tafasitamab-cxix.

Fertility studies have not been conducted with tafasitamab-cxix.

In the 13-week repeat-dose general toxicity study in cynomolgus monkeys, no adverse effects on male and female reproductive organs were observed up to the highest dose tested, 100 mg/kg/week (approximately 9 times the human exposure based on AUC at the clinical dose of 12 mg/kg/week).

## 14 CLINICAL STUDIES

The efficacy of MONJUVI in combination with lenalidomide followed by MONJUVI as monotherapy was evaluated in L-MIND, an open-label, multicenter, single arm trial (NCT02399085). Eligible patients had relapsed or refractory DLBCL after 1 to 3 prior systemic therapies, including a CD20-directed cytolytic antibody, and were not candidates for high dose chemotherapy (HDC) followed by autologous stem cell transplantation (ASCT). Patients received MONJUVI 12 mg/kg intravenously in combination with lenalidomide (25 mg orally on Days 1 to 21 of each 28-day cycle) for a maximum of 12 cycles, followed by MONJUVI as monotherapy until disease progression or unacceptable toxicity as follows:

- Cycle 1: Days 1, 4, 8, 15 and 22 of the 28-day cycle;
- Cycles 2 and 3: Days 1, 8, 15 and 22 of each 28-day cycle;
- Cycles 4 and beyond: Days 1 and 15 of each 28-day cycle.

Of the 71 patients with DLBCL confirmed by central laboratory who received the combination therapy, the median age was 71 years (range: 41 to 86 years); 55% were males, and 100% had received a prior CD20-containing therapy. Race was collected in 92% of patients; of these, 95% were White, and 3% were Asian. The median number of prior therapies was two; 49% had one prior line of treatment, and 51% had 2 to 4 prior lines. Thirty-two patients (45%) were refractory to their last prior therapy and 30 (42%) were refractory to rituximab. Nine patients (13%) had received prior ASCT. The primary reasons patients were not candidates for ASCT included age (47%), refractoriness to salvage chemotherapy (27%), comorbidities (13%) and refusal of high dose chemotherapy/ASCT (13%).

Efficacy was established based on best overall response rate, defined as the proportion of complete and partial responders, and duration of response, as assessed by an Independent Review Committee using the International Working Group Response Criteria (Cheson 2007). Results are summarized in **Table 5**.

**TABLE 5: Efficacy Results in L-MIND**

	<b>N = 71</b>
<b>Best overall response rate, n (%)</b>	39 (55%)
(95% CI)	(43%, 67%)
Complete response rate	37%
Partial response rate	18%
<b>Duration of Response</b>	
Median (range) in months <sup>a</sup>	21.7 (0, 24)

<sup>a</sup>Kaplan Meier estimates

## 16 HOW SUPPLIED/STORAGE AND HANDLING

MONJUVI (tafasitamab-cxix) for injection is a sterile, preservative-free, white to slightly yellowish lyophilized powder for reconstitution supplied as a 200 mg single-dose vial.

Each 200 mg vial is individually packaged in a carton (NDC 73535–208–01).

Store refrigerated at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. Do not shake. Do not freeze.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

### Infusion-Related Reactions

Advise patients to contact their healthcare provider if they experience signs and symptoms of infusion-related reactions [see *Warnings and Precautions (5.1)*].

### Myelosuppression

Inform patients about the risk of myelosuppression. Advise patients to immediately contact their healthcare provider for a fever of 100.4°F (38°C) or greater or signs or symptoms of bruising or bleeding. Advise patients of the need for periodic monitoring of blood counts [see *Warnings and Precautions (5.2)*].

### Infections

Inform patients about the risk of infections. Advise patients to immediately contact their healthcare provider for a fever of 100.4°F (38°C) or greater or signs or symptoms of infection [see *Warnings and Precautions (5.3)*].

### Embryo-Fetal Toxicity

- Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to inform their healthcare provider of a known or suspected pregnancy [see *Warnings and Precautions (5.4)*, *Use in Specific Population (8.1)*].
- Advise females of reproductive potential to use effective contraception during treatment with MONJUVI and for at least 3 months after the last dose [see *Use in Specific Populations (8.3)*].
- Advise patients that lenalidomide has the potential to cause fetal harm and has specific requirements regarding contraception, pregnancy testing, blood and sperm donation, and transmission in sperm. Lenalidomide is only available through a REMS program [see *Use in Specific Populations (8.1,8.3)*].

### Lactation

Advise women not to breastfeed during treatment with MONJUVI and for at least 3 months after the last dose [see *Use in Specific Populations (8.2)*].

Manufactured by:  
**MORPHOSYS US INC.**  
Boston, MA 02210  
U.S. License No. 2512

Marketed by:  
**MORPHOSYS US INC. and INCYTE Corporation.**  
MONJUVI is a registered trademark of MorphoSys AG

Copyright © 2020 MorphoSys AG. All rights reserved.  
Product of Germany



MONJUVI and the MONJUVI logo are registered trademarks of MorphoSys AG.  
© 2020

November 2020 RC-US-TAF-00525

Distributed and marketed by MorphoSys US Inc. and marketed by Incyte Corporation  
MorphoSys is a registered trademark of MorphoSys AG.  
Incyte and the Incyte logo are registered trademarks of Incyte Corp.